


Medicines for high blood pressure

Patient information: medicines




Working together for better patient information

What is this leaflet about?

High blood pressure is common in people with kidney disease. Most people with high blood pressure will eventually need medicines to treat it, which also helps protect your kidneys from further damage. People with kidney disease are at a higher risk of having heart attacks and strokes. One of the best treatments for this is to treat high blood pressure.

This leaflet aims to give you more information about your blood pressure medicines and some of their possible side effects. For more information, ask your doctor or pharmacist.





Why do I need to take different types of medicines for high blood pressure?

There are many medicines available to reduce blood pressure. Your doctor will have chosen those most suitable for you. A combination of a few medicines is often needed to reach your target blood pressure.

High blood pressure may not make you feel unwell, but it can cause other health problems and damage your kidneys, so it is important to continue taking your medicines even if you don't feel any different. Talk to your doctor if you have any side effects.



What types of medicines might I need?

Some commonly used medicines for blood pressure are listed in the table. They will be prescribed to you as tablets. Speak to your doctor or pharmacist if you are taking a medicine that is not listed here.

Blood pressure medicine	Example of drugs in this group	Common side effects
Alpha-blockers	Doxazosin	Dizziness, headache, incontinence in women
ACE-inhibitors or 'prils'	Lisinopril, perindopril, ramipril	Cough
ARBs or 'sartans'	Losartan, candesartan, irbesartan, valsartan	Tiredness
Beta-blockers	Atenolol, bisoprolol, propranolol	Cold hands and feet, tiredness
Calcium antagonists	Amlodipine, felodipine, nifedipine, diltiazem, verapamil	Swollen ankles
Other drugs	Moxonidine, hydralazine	Tiredness and a build up of fluid in the body with hydralazine
Water tablets (diuretics)	Furosemide, bendroflumethiazide, spironolactone	Dehydration, dizziness, increase or decrease in potassium levels in the blood.



I'm on different medicines from those in the table, is that ok?

Yes. There are many different types of blood pressure medicines.

Although some are more commonly prescribed, sometimes there are special reasons to use different medicines, particularly if your blood pressure is difficult to control, there are special circumstances such as pregnancy, or you have had side effects with the medicines.

I feel dizzy, is that a side effect?

Like any medicine, blood pressure tablets can have side effects.

Many of these side effects are not dangerous and often get better with time. If you do have side effects it is important not to stop all of your medicines suddenly as this can lead to your blood pressure increasing. Make sure that you talk to your doctor or nurse to advise you.

A common effect of blood pressure tablets is feeling dizzy when you stand up. This can last a few seconds or longer. It is helpful to try not to stand up too quickly and make sure you have a blood pressure check to keep an eye on your blood pressure as sometimes changes in the type of medicine and dose can help.

My doctor mentioned 'Sick Day Rules'. What does that mean?

Taking certain medicines when you are unwell can lead to more severe illness. 'Sick Day Rules' is advice about how to take your medicine if you are unwell and become dehydrated.

Your doctor or nurse may suggest that you stop some of your medicines. Dehydration occurs if you are unable to drink fluid or have diarrhoea or vomiting. You should stay off these medicines until you are eating and drinking normally for at least 48 hours.

Examples of blood pressure medicines you may be advised to stop:

- ACE-inhibitors
- ARBs
- Diuretics ('water tablets')

Other medicines you may be advised to also stop taking:

- Metformin
- Tolvaptan
- SGLT-2 inhibitors ('flozins')

How should I take my blood pressure medicines with haemodialysis?

Sometimes blood pressure medicines can cause your blood pressure to fall too low during haemodialysis.

If this happens, the type of blood pressure medicine you are prescribed may be changed, or you may be advised to change the time when you take your medicines, for example, taking them at night if you have dialysis in the morning.



Looking after your medicines

You should store your medicines in their original packet in a cool, dry place, out of the sight and reach of children.

Do not get rid of any expired or unwanted tablets by flushing them down the toilet, or throwing them away. Take them to your local pharmacy who will dispose of them for you.

Where can I find out more information?

- **Kidney Care UK** Medication - www.kidneycareuk.org/medication
- **Kidney Care UK** high blood pressure - www.kidneycareuk.org/high-blood-pressure
- **Kidney Care UK** Blood Pressure Association - www.bloodpressureuk.org
- **NHS** Medicines A to Z - www.nhs.uk/medicines

Contact us to see how we can support you

Call free on
0808 801 00 00

(Monday to Friday, 9am-5pm)

support@kidneycareuk.org

www.kidneycareuk.org



If you have feedback about this leaflet, please let us know at: feedback@kidneycareuk.org

Publication date 07/2023
Review date 07/2026



in partnership with

UK Kidney Association

© Kidney Care UK. Kidney Care UK is a charitable company limited by guarantee. Registered in England and Wales (1228114). A charity registered in England and Wales (270288) and Scotland (SCO48198). KC65.